



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A Proof of the Theorem


THE EQUATION $f(z) = 0$ HAS A ROOT WHERE $f(z)$ IS ANY HOLOMORPHIC FUNCTION OF z .

BY J. C. FIELDS, *Fellow in Mathematics, Johns Hopkins University.*

Represent z by a point in one plane and $f(z) = \zeta$ by a point in another plane.

If $f(z) = \zeta$ cannot become equal to zero for any value of z , there must be some minimum distance from the origin ω within which ζ cannot fall. Suppose ζ at such minimum distance from ω .

We can suppose this, for, $f(z)$ becoming infinite along with z , this minimum must be for a finite value of z , and can therefore be reached.



To z give an increment δz ; the corresponding increment of ζ is $\delta \zeta = \frac{f^r(z)}{r} (\delta z)^r$ where $f^r(z)$ is the first of the successive derivatives of $f(z)$ which does not vanish for this value of z . Now varying δz , make it describe a closed curve round z ; $\delta \zeta$ will at the same time describe a closed curve r times over round the point ζ , and will therefore come between the point ζ and origin ω . The point ζ has then no such minimum distance from the origin ω as was supposed; the function $f(z) = \zeta$ is therefore capable of becoming $= 0$, and the equation $f(z) = 0$ has therefore a root.

The above statement might be slightly varied, thus: to $f(z)$ we can give an increment in any direction we may choose; for δz , $\delta \zeta$, being any two corresponding increments of z , ζ , respectively; if the required increment is to be in a direction inclined to $\delta \zeta$ at an angle α , give to δz the rotation $\frac{\alpha}{r}$ and $\delta \zeta$ takes the required direction.

If mod (δz) remain constant while rotating round z , $\delta \zeta$ at the same time describes a circle round ζ , and we give ζ as much of an increment in one direction as in another.

We can always for a polynomial take $r=1$, for if there be no root of $f(z)=0$, ζ will have a minimum along each of radiating lines drawn through origin ω (for $f(z)$ can easily be shown to have some value on each of these lines), and as $f'(z)$ cannot vanish for more than $n-1$ values of z , we can always choose our minimum along a line on which $f'(z)$ cannot vanish.

[Just as this note was about to go to press, I discovered that practically the same proof as above had been given by Hoüel in his Cours de Calcul Infinitésimal.—J. C. F.]